


Cervical Facet/Medial Branch Block (MBB)


What is a facet joint pain?

The spine is made of vertebrae, which makes up the spine. The vertebrae are connected to each other with facet joints, which allows the bending and rotational spine movements. As the joints become inflamed and irritated, there is a small medial branch nerve that transmits the pain signal from the joint to the brain. Furthermore, the pain may worsen when the spine is extended.



What are the indications of cervical medial branch block?

Among many indications the listed are the main indications: spondylosis, spondylolisthesis, arthritis, osteoarthritis, spondyloarthritis, chronic headaches, neck pain, failed neck surgery syndrome.



How does cervical medial branch block bring pain relief?

The pain is produced due to inflammation cervical facet joint which is transmitted via medial branch nerve to the central nervous system. By injection local anesthetic which helps numbs out the medial branch nerve, this results in decreased neck pain caused by the facet joints.

What are risks for the cervical medial branch block?

Among many, here are few listed: increased pain, infection, bleeding, nerve damage, weakness, numbness, muscle spasms, headache.

How is the cervical medial branch block performed?

After sterile preparation of the neck region, the injection site is localized under X-ray. Following the local anesthetic applied to the injection site, which can help decrease the injection site pain, the needle is guided toward the target cervical facet joint with the help of X-ray. After the target is reached, it can be further confirmed with liquid contrast. Thereafter, small amount of local anesthetic is injected, and the needle is taken out at the end of the procedure.

What to expect after the procedure?

This is an **outpatient procedure**. Patient should expect to receive instant >80% relief from the joint nerve blocks.

How long the relief from the cervical MBB would last for?

It varies from patient to patient. Usually, the pain relief can last for few hours to few months.

What is the purpose of the cervical medial branch block?

Cervical MBB is a diagnostic procedure **NOT** therapeutic procedure. It helps confirm the pain arising from the cervical joints. If patient receives > 80% relief from two separate cervical medial branch block, then the patient may be an excellent candidate for cervical radiofrequency ablation of the medial branch nerves which can bring 6-18 months of pain relief. Please read the cervical radiofrequency handout for more details.

Please tell your pain physician if you are taking any blood thinners. For example: Coumadin, Plavix, Heparin, Lovenox and etc. The blood thinners need to be stopped before the interventional procedure, and the time frame will be prescribed by your pain physician.